

2. Obstruction of small right coronary arteries that predominantly supply the right ventricle when the inferior wall of the left ventricle is supplied by a dominant left circumflex artery.

3. Obstruction of a right ventricular branch of a dominant right coronary artery or occlusion of a dominant right coronary artery proximal to the right ventricular branch. In the last two variants, we observed concomitant ST elevation in inferior and anterior leads. In the 2nd variant (obstruction of a small right coronary artery), we observed isolated right ventricular infarction (2) and, in the third, infarction of the right ventricle and of the inferior wall of the left ventricle.

These data suggest that concomitant ST elevation in inferior and anterior leads cannot differentiate between patients with single or multivessel disease and in all cases indicates an extensive ischemic process. We believe in mandatory early catheterization for every patient with such concomitant ST elevation.

ILYA OVSYSHCHER, MD, PhD  
Ben-Gurion University  
P.O. Box 151  
Beer-Sheva, Israel

#### References

1. Lew AS, Hod H, Cereck R, et al. Inferior ST segment changes during acute anterior myocardial infarction: a marker of the presence or absence of concomitant inferior wall ischemia. *J Am Coll Cardiol* 1987;10:519-26.
2. Ila R, Margulies G, Goldfarb B, et al. ST elevation in leads V<sub>1</sub> to V<sub>4</sub> caused by isolated right ventricular infarction and infarction. *Cardiology* 1987;74:598-9.

#### Reply

We thank Ovysysher for emphasizing the earlier observations of our group (1,2) and others (3) that combined anterior and inferior lead ST segment elevation may be due to inferior infarction with predominant or extensive right ventricular involvement. We agree that the electrocardiographic (ECG) injury pattern during acute myocardial infarction generally reflects the anatomic supply of the artery of infarction rather than the extent of coronary artery disease (4), although these two factors may be interactive in the setting of extensive coronary artery disease with well developed collateral perfusion.

In general, we also agree with Ovysysher's recommendation that patients with concomitant anterior and inferior lead ST segment elevation be considered for early coronary angiography (as a prelude to reperfusion of the ischemic myocardium), but we do not believe that the ECG pattern in itself mandates this approach because the decision must also be based on other determinants of patient suitability (duration of symptoms, for example).

ALLAN S. LEW, MD  
Cedars-Sinai Medical Center  
Los Angeles, California 90048

#### References

1. Grif HL, Shih PK, Rodriguez L, et al. ST elevations in leads V<sub>1</sub> to V<sub>4</sub> may be caused by right coronary artery occlusion and acute right ventricular infarction. *Am J Cardiol* 1984;53:991-6.
2. Lew AS, Madhavi J, Shih PK, Weiss AT, Peter T, Ganz W. Factors that determine the direction and magnitude of precordial ST segment changes during acute inferior myocardial infarction. *Am J Cardiol* 1985;55:883-8.
3. Lewis BS, Rosenman Y, Ben-David Y. The hyperacute phase of right ventricular infarction. *Heart Lung* 1984;13:683-4.
4. Lew AS, Weiss AT, Shih PK, et al. Precordial ST segment depression during acute inferior myocardial infarction: early thallium-201 scintigraphic evidence of adjacent posterolateral and/or inferoseptal involvement. *J Am Coll Cardiol* 1985;5:304-9.

### Electrocardiogram in Right Ventricular Infarction

In his lucid editorial on the new "risk factor" for right ventricular infarction (1), Krasnow seems to introduce new electrocardiographic criteria for that condition. The "prominent though transient R waves in the right precordial leads of the electrocardiogram, especially lead V<sub>1</sub>R" have not, to my knowledge, been described as markers of right ventricular infarction. The morphologic features described are those of (true) posterior infarction, which may affect the diagnosis of right ventricular infarction by making the ST segment in the right precordial leads isoelectric (2). I feel that Krasnow may be in error, perhaps referring to the R waves of right ventricular hypertrophy in those leads?

GEORGE NIKOLIC, MBBS, FRACP, FACC  
Woden Valley Hospital  
P.O. Box 11, Woden  
Australian Capital Territory 2606, Australia

#### References

1. Krasnow N. Right ventricular hypertrophy: a new "risk factor" for right ventricular infarction. *J Am Coll Cardiol* 1987;10:1188-9.
2. Reddy GV, Schamrahn L. The electrocardiology of right ventricular myocardial infarction. *Chest* 1986;90:756-60.

#### Reply

Nikolic is correct in that the transient electrocardiographic changes of right ventricular infarction should have referred to ST change and not R waves. I was indeed thinking of right ventricular hypertrophy, the subject under discussion. I thank him for his correction and the care with which he read the editorial.

NORMAN KRASNOW, MD, FACC  
State University of New York  
Health Science Center at Brooklyn  
450 Clarkson Avenue  
Brooklyn, New York 11203